



Figure 4: Cellulitis of the left arm following use of the proximal AVF

SHORTAGE OF SKILLED MANPOWER

For a country of 1.2 billion, there are only about 1000 trained nephrologists. This is obviously way short of the numbers actually needed. Besides, there is immense shortage of skilled dialysis technicians / nurses in the country. Till recently there was no recognized course of for training this cadre. To overcome this problem, the Govt of India is starting courses specifically to train them.

INACCESSIBILITY OF HEMODIALYSIS CENTER

India is a vast country. More than 70% of the population lives in villages. Hemodialysis centres are present only in the cities and large towns. Until recently several district headquarters did not have HD facilities. Hence it is logistically difficult for the vast majority of dialysis patients to undergo regular maintenance hemodialysis (MHD) on a long term basis.

ECONOMICS OF HEMODIALYSIS

The number of patients on renal replacement therapy (RRT) in general, and hemodialysis in particular reflects the socioeconomic condition of the country. 26% of the 1.2 billion population subsists below the poverty line, where the daily earning is ₹ 10. The government spends barely ₹ 400 per capita on health. The priorities are on infectious diseases, sanitation and nutrition. Maintenance hemodialysis is not provided in government hospitals. Insurance covers a tiny fraction of the population. Most patients are self-paying. The monthly cost of HD in most private hospitals is about ₹ 12000, added to which is the cost of medications.⁷ This amount is beyond the reach of the vast majority. In cases where the patient lives far away from the dialysis centre it entails further expenses of travel and wages lost for the day. In the NIMS study at the end of the follow-up period the number of self-paying patients had dwindled significantly. Hence underdialysis with all its attendant complications is the norm. While the science of hemodialysis dictates stringent adequacy measures with mounting evidence of the advantages of daily dialysis, many patients survive on the edge by taking dialysis once a week. Few years back the concept of stand-alone HD units on public-private partnership was conceived to provide cheaper dialysis. Such units are gradually becoming functional all over the country. The involvement of philanthropists and non-governmental organizations in the setting up of charitable dialysis units has also helped in providing affordable dialysis to needy patients.⁷ These stand-alone dialysis units need to be associated with a proper hospital, so that if needed unstable patients can be shifted for admission.

MALNUTRITION

Given the economics and logistics of hemodialysis, it is not surprising that malnutrition is very common in MHD patients. In a study from South India, 91% of the patients were moderately malnourished.⁸ The role of renal dieticians is of great importance to remedy this condition.⁹ Unfortunately there is a dearth of committed renal dieticians in the country. It is well known that malnutrition is usually associated with the Malnutrition-Inflammation-Atherosclerosis Syndrome, which leads to an overall poor outcome due to sepsis and cardiovascular mortality.

HEPATITIS SEROCONVERSION DURING MHD

Acquiring Hepatitis B and or C infection while on dialysis is another major problem in the country. Availability of reasonably effective vaccination against hepatitis B has helped to reduce the conversion rate. However the scenario is dismal with regard to hepatitis C. There is no vaccination to prevent hepatitis C, and the spread of the virus is both blood-borne and nosocomial. The prevalence is widely variable, from <5% to 75%.¹⁰⁻¹¹ The prevalence is consistently associated with the number of transfusions, and more importantly the duration of MHD. A study by Jasuja et al from a tertiary care hospital in Delhi showed that almost half the patients with duration of dialysis more than 16 months were HCV RNA positive.¹² The CDC guidelines do not recommend isolation of hepatitis C positive cases during dialysis, provided universal precautions (UP) are meticulously adhered to. Implementation of UP requires knowledge about it, commitment to adhere to it and additional costs. A study at our centre showed that though all the nurses were aware about them, only 70% followed them in day-to-day practice.¹³ Hence it is our opinion that if a unit is to contain HCV infection, then they must isolate HCV-positive patients ie. the 'no-isolation' policy should not be generalized.¹⁴

CONCLUSION

Hemodialysis in India has improved by leaps over the last two decades. However there is a long way to go. Intensive public, patient and primary care physician education, government backing and induction of more trained personnel will make hemodialysis a successful RRT modality in India.

BIBLIOGRAPHY

1. Chugh KS. Five decades of Indian nephrology: A personal journey. *Am J Kidney Dis* 2009;54:753-763.
2. Swarnalatha G, Ram R, Prasad N, Dakshinamurthy KV. End-stage renal disease patients on hemodialysis: A study from a tertiary care centre in a developing country. *Hemodial Int* 2011;15:312-319.
3. Kher V. Tunneled central venous catheters for dialysis - A necessary evil? *Indian J Nephrol* 2011;21:221-222.
4. Bhowmik D, Baveja T, Tiwari SC. Role of the family physician in controlling the epidemic of chronic kidney disease. *J Indian Med Assoc* 2012;110:40-42.
5. KDOQI Clinical Practice Guidelines and Clinical Practice Recommendations 2006 Updates Hemodialysis adequacy Peritoneal Dialysis Adequacy Vascular Access. *Am J Kidney Dis* 2006;48(Suppl 1):S1.
6. Srivastava A, Sharma S. Hemodialysis vascular access options after failed Brescia-Cimino arteriovenous fistula. *Indian J Urol* 2011;27:163-167.
7. Khanna U. The economics of dialysis in India. *Indian J Nephrol* 2009;19:1-4.
8. Janardhan V, Soundararajan P, Vanitha Rami N, et al. Prediction of malnutrition using modified subjective global assessment-dialysis malnutrition score in patients on hemodialysis. *Indian J Pharm Sci* 2011;73:38-45.
9. Colman S, Bross R, Brenner D. *J Ren Nutr* 2005;15:231-43. The Nutritional and Inflammatory Evaluation in Dialysis patients (NIED) study: overview of the NIED study and the role of dietitians.
10. Pol S, Vallet-Pichard A, Fontaine H, Lebray P. HCV infection and hemodialysis. *Semin Nephrol* 2002;22:331-339.
11. Covic A, Lancu L, Apetrei C, et al. Hepatitis virus infection in hemodialysis patients in Moldavia. *Nephrol Dial Transplant* 1999;14:40-45.
12. Jasuja S, Gupta AK, Choudhry R, et al. Prevalence and associations of hepatitis C viremia in hemodialysis patients at a tertiary care hospital. *Indian J Nephrol* 2009;19:62-67.
13. Agarwal SK, Mohan MP, Varghese M. Assessment of awareness regarding universal precaution among the nursing staff of AIIMS in 1997. *J Assoc Physicians India* 1998;46:1061.
14. Agarwal SK, Dash SC, Gupta S, Pandey RM. Hepatitis C virus infection in hemodialysis: The 'no-isolation' policy should not be generalized. *Nephron Clin Pract* 2009;111:c133-c140.

Future Special Issues/ Symposia

Special Issues

- Recent Advances & Future Trends in Healthcare
- Imaging in the 21st Century
- Advances in Pediatric Surgery
- HIV/AIDS: Emerging Trends

Symposia

- Drug Addiction: Health Hazards
- Sleep Disorders: Current Perspective